

Effects of two weekly servings of cod for 16 weeks in pregnancy on maternal iodine status and infant neurodevelopment: Mommy's Food, a randomized controlled trial

Maria Wik Markhus, Mari Hysing, Lisa Kolden Midtbø, Ive Nerhus, Synnøve Næss, Inger Aakre, Ingrid Kvestad, Lisbeth Dahl, and Marian Kjellevoid.

Maria Wik Markhus, PhD (maria.wik.markhus@hi.no)^{1*}, Mari Hysing, PhD (mari.hysing@uib.no)^{2,3}, Lisa Kolden Midtbø, PhD (lisa.kolden.midtbo@moreforsk.no)¹, Ive Nerhus, MSc (i.nerhus@gmail.com)¹, Synnøve Næss, MSc (synnoeve.naess@hi.no)¹, Inger Aakre, PhD (Inger.Aakre@hi.no)¹, Ingrid Kvestad, PhD (inkv@norceresearch.no)², Lisbeth Dahl, Dr. Scient. (lisbeth.dahl@hi.no)¹, and Marian Kjellevoid, Dr. Scient. (marian.kjellevoid@hi.no)¹

Author affiliations

¹Seafood, Nutrition and Environmental State, Institute of Marine Research, P.O box 1870 Nordnes, NO-5817 Bergen, Norway

²Regional Centre for Child and Youth Mental Health and Child Welfare, NORCE Norwegian Research Centre, P.O box 22 Nygårdstangen, NO-5838 Bergen, Norway

³Department of Psychosocial Science, Faculty of Psychology, University of Bergen, P.O box 7807, 5020 Bergen, Norway

*Corresponding author

Keywords: Iodine; Thyroid hormones; Pregnancy; Neurodevelopment; RCT; Infants

Running title: Maternal iodine status trial and infant neurodevelopment

Abstract

Background

Mild-to-moderate iodine deficiency is still present in many countries, particularly in pregnant women. Observational studies suggest that mild-to-moderate iodine deficiency during pregnancy may be associated with impaired thyroid function and child neurodevelopment. Randomized controlled food trials to increase iodine status are scarce. We assessed the impact of an increased intake of cod during pregnancy on maternal iodine status, and infant neurodevelopment.

Methods

In this randomized controlled trial, pregnant women in Bergen, Norway, recruited through Haukeland University Hospital, were randomly assigned (1:1) to an intervention of 200 g of cod twice a week for 16 weeks (gestational week 20-36) or to continue with their standard diet (control group). Randomization was done by lottery. Primary outcome was urinary iodine concentration (UIC) (spot samples from six consecutive days) measured post intervention. Secondary outcome was infant neurodevelopment assessed by the cognitive, language and motor scales of the Bayley Scales of Infant and Toddler Developmental 3rd edition (Bayley-III) at 11 months of age. In addition, maternal thyroid function was measured (TSH, fT3, fT4) at baseline and post intervention. The trial was registered in ClinicalTrials.gov, NCT02610959.

Results

Between Jan 2016 until Feb 2017, 137 women were recruited. Post intervention UIC was higher in the intervention group (n=61) (median (IQR) 98 (64-145) $\mu\text{g/L}$), compared to control (n= 61) (median (IQR) 73 (52-120) $\mu\text{g/L}$) ($p= 0.028$), also after adjusting for baseline UIC ($p=0.048$). Infants of mothers in the intervention group had a lower cognitive composite score on the Bayley-III compared to the control group, ($p= 0.045$). There were no group differences in the Bayley III language- or motor composite scores. Maternal thyroid hormones (TSH, fT3, fT4) did not differ between the groups post intervention.

Conclusions

Increased cod intake during pregnancy improved the iodine status in women with mild-to-moderate iodine deficiency, however, did not affect thyroid function. The negative effect on cognition should be followed up to assess whether this is a stable effect over time. More studies are warranted to enable good health advice on iodine nutrition in pregnancy.

Trial registration

ClinicalTrials.gov NCT02610959. Registered November 20, 2015.

Introduction

Although substantial progress has been made towards improving iodine status worldwide, iodine deficiency remains a significant health problem both in low-, middle- and high-income countries (1). Severe iodine deficiency is almost completely eradicated, mainly through salt iodization programs (2). However, mild-to-moderate iodine deficiency is still present in many countries, particularly in women of childbearing age, where Europe is the continent with the highest prevalence of iodine deficiency worldwide (3).

Iodine is an essential micronutrient because of its incorporation in the thyroid hormones triiodothyronine (T3) and thyroxine (T4). Pregnancy is a period of increased iodine requirements in order to maintain euthyroidism due to transfer of iodine and thyroid hormones to the fetus (4). Consequently, pregnant women are vulnerable to iodine deficiency.

The World Health Organization (WHO) recommends an iodine intake of 150 µg/day for non-pregnant women and 250 µg/day for pregnant women and the Nordic Nutrition Recommendations recommend 150 µg/day for non-pregnant women, 175 µg/day for pregnant women, and 200 µg/day for lactating women (2, 5). WHO recommends using the median urinary iodine concentration (UIC) to assess iodine status in population groups. A median UIC of 150-250 µg/L indicates optimal iodine nutrition in pregnant women, while a UIC <150 is considered insufficient (2). The Nordic Nutrition Recommendations of 175 µg/day corresponds to a UIC of ~100 µg/L. Observational studies have found adverse health effects when UIC falls below 100 µg/L, indicating that this may be proposed as a cut-off for sufficiency during pregnancy. (6, 7). Norway is currently a country of documented mild-to-moderate iodine deficiency in pregnant women with median UIC ranging from 75-92 µg/L (8-11). This may be caused by the decrease in intake of milk and lean fish during the last decade's (12). Norway has no iodized salt, and milk, dairy products and seafood are the main dietary iodine sources of the total iodine intake (8, 13). After the completion of this trial, Norwegian authorities have recommended iodine supplementation of 150 µg/day to pregnant women whom have a lower daily intake than

6 dl of cow's milk / yogurt (but eat white saltwater fish regularly), or eat little / no white saltwater fish and at the same time have a lower daily intake than 8 dl cow's milk / yogurt.

Severe iodine deficiency is a well-known risk factor for cognitive deficits in children (14, 15). The effects of mild-to-moderate iodine deficiency on cognitive development is less certain. Few intervention studies on maternal iodine supplementation on child development exist and randomized control trials (RCTs) have been urgently called for (16, 17). To the best of our knowledge, this is the first intervention trial in pregnant women with iodine rich food (cod) to study its effects on maternal iodine status and infant neurodevelopment. The primary aim of this RCT was to investigate if an increased intake of cod during pregnancy has an effect on maternal iodine status, and secondary if it has an effect on infant neurodevelopment at 11 months of age.

Materials and methods

Study design and participants

This study was a two-arm RCT with a primary aim of studying the effect of increased cod intake during pregnancy on maternal iodine status and secondary aim of determining infant neurodevelopment when the children were 11 months old. Participants were recruited through the Women's Clinic at Haukeland University Hospital, Health region West, Norway. Approximately 5000 women give birth at the Women's Clinic annually. Once a pregnant woman was enrolled, the study investigators made every reasonable effort to follow the participant closely for the entire trial period in order to ensure the best possible retention. Prior to all the visits, the participants were reminded about the upcoming appointment.

From January 2016 until February 2017, information regarding the intervention trial was included in the invitation from the Women's Clinic at the time of routine ultrasound in gestational week 18. To increase the enrolment rate, information regarding the trial, and invitation to participate was also broadcasted online (Facebook, Instagram, and in an online magazine for pregnant women in Norway). Pregnant women who were interested in the study contacted the researchers at the Institute of Marine Research (IMR), Bergen, Norway. Inclusion criteria were primiparous singleton pregnancy,

gestational week \leq 19, Norwegian speaking and/or understand Norwegian writing. Exclusion criteria were allergies to fish and chronic diseases known to affect iodine status (Graves' disease, thyroiditis, thyroid nodules, known hypothyroidism or hyperthyroidism). Participants gave written informed consent after receiving written and oral information about the study. No specific information about iodine was given as the overall aim of the study was to study the relationship between Mommy's Food (nutrient intake in pregnancy and the infants first year) and infant development. The women could withdraw from the study at any time without giving any reason. The trial complies with the Declaration of Helsinki and was approved by the Regional Committee for Medical and Health Research Ethics West (2015/879). The study protocol has been published elsewhere (18).

Randomization and masking

Baseline data was collected at the first visit, and during gestational week 18-19. At the second visit in gestational week 19, the participants were randomized individually by lottery in blocks of ten to ensure approximately equal allocation to both groups. Owing to the nature of the intervention, blinding of the participating mothers was not possible. Study investigators (LKM and IN) generated the random allocation sequence, enrolled participants, and assigned participants to groups. Laboratory personnel were blinded when analyzing data with no access to the code. Study investigators (MWM and SN) were blinded when analyzing the primary and secondary outcomes as the code was masked with dummy variables by a third study investigator (LKM).

Intervention

After randomization, participants in the intervention group received frozen cod fillets (Lerøy A/S, Bergen, Norway) and were instructed to consume two intervention meals of 200 g weekly (a total of 400 g cod per week) for 16 weeks (a total of 32 meals) from gestational week 20 to 36. The participants also received cod for their partner, if any, with the intention to increase compliance. The participants prepared the meals themselves and could choose their own recipes but were also provided with recipes that could be used ad libitum. For compliance purposes, participants were instructed to weigh (Kitchen Scale, article no. 34-1207-16, ClasOhlson.com) the cod fillet before preparing the meals, and

weigh the fillet leftovers (if any), after the meals were eaten. The participants recorded these data in a weight registration form, in addition to recipes used, and date of consumption. To calculate total g cod eaten each meal, g cod after preparation of the meal were subtracted from g cod before preparing the meal. Total g cod for each meal during the intervention period was then summarized to get a value of total g cod eaten during the intervention period. Total intake of cod was divided by 16 weeks to get a mean weekly intake of cod. Total intake of cod was divided by maximum intake of cod during the intervention period (200 g x 32 meals = 6400 g) and multiplied with 100 to get a compliance score. Example a total intake of 6400 g during the intervention or a mean intake of 400 g cod per week provided a compliance score of 100. The participants in the control group were instructed to continue to follow their habitual diet, without any restrictions.

Outcomes

The primary outcome was UIC measured at post intervention. At the first visit in gestational week 18 participants received six marked collection tubes for the collection of urine samples on six consecutive days from gestational week 18-19, with instructions on how to collect the spot urine samples. Prior to the visit in gestational week 36, post intervention, participants received by mail six marked collection tubes for the collection of urine samples on six consecutive days from gestational week 35-36. The participants kept the urine samples in their home freezer until the visits in gestational week 19 and 36, respectively. Equal amounts of urine from the six spot urine samples (collected on six consecutive days between 4 pm and midnight) were homogenized into one pooled sample of 1 ml urine and were stored at minus 20°C in cryo tubes (CryoTube™ Vials Nunc, Thermo Fischer Scientific, Roskilde, Denmark) pending analysis by inductively coupled plasma mass spectrometry (ICP-MS). Prior to the analysis, the urine samples were defrosted in a refrigerator, diluted with 1% tetramethylammonium hydroxid (TMAH), filtrated using a sterile membrane filter (0.45 µm pore size) and transferred to tubes appropriate for the analysis by the Agilent 7500 for ICP-MS at IMR. Samples were analyzed against a urine calibration curve (standard addition curve) to measure the unknown iodine concentration (127I) in the collected urine samples. Accuracy was verified with certified reference

material; Seronorm Trace Elements Urine (Nycomed Pharma, Norway), iodine content: 84 µg/L (range, 72-96 µg/L) and 304 µg/L (range: 260-348 µg/L). The measurement uncertainty of the method has been assessed based on internal reproducibility, analysis of standard reference material (SRM) and is set at 20% in the entire range (2-297 µg/L). The limit of detection was 2 µg/L.

For thyroid function and docosahexaenoic acid (DHA) analysis blood samples were drawn in gestational week 18 and 36. Blood samples for serum preparation were collected in BD Vacutainer® SST™ vials II *Advanced* and set to coagulate for minimum 30 min before centrifuging (1000-3000 G, room temperature, 10 min) within 60 minutes after extraction. Blood samples for red blood cell preparation were collected in BD Vacutainer® K2E 5.4 mg vials and centrifuged (1000–1300 G, 20 °C, 10 min) within 30 min. Post separation, serum samples were stored at minus 80 °C pending analysis at Fürst Medical Laboratories, Norway, and red blood cell samples were stored at minus 80 °C pending analysis at IMR. The serum samples were stored for a maximum of three months before analysis. Thyroid stimulating hormone (TSH), free T4 (fT4) and free T3 (fT3) were analysed in serum using magnetic separation and detection by chemiluminescence, labelled with acridinium ester, on an Advia Centaur XPT Immunoassay system (Siemens Healthcare diagnostics Inc., Tarrytown, NY, USA). For all blood constitutes the coefficient of variation (CV) was <6%. DHA was analysed in red blood cells by a standardized procedures at IMR (19), using ultrafast gas chromatographic (Thermo Electron Corporation, Franklin, MA, USA).

The secondary outcome was neurodevelopment assessed by the cognitive, language and motor scales of the Bayley Scales of Infant and Toddler Developmental 3rd edition (Bayley-III) when the infants were 11 months of age. The Bayley-III is a comprehensive assessment tool of neurodevelopment administered directly with the child (20). The tool takes approximately 45 to 60 minutes to administer and includes three main subscales: the cognitive, language (receptive and expressive) and motor (fine and gross motor) scales. The Bayley-III represents the gold standard of developmental assessment in this age group and is widely used as an outcome measure in clinical trials. The official Norwegian version of the Bayley-III translated and adapted for a Norwegian setting was used, with American norms from a representative American sample. The Bayley-III

provides 5 scaled scores (mean 10, range 1-19, standard deviation (SD) 3) and 3 composite scores (mean 100, range from 40-160 and, SD of 15) (21). Two trained testers (LKM and IN), supervised by a neuropsychologist (MH) and a clinical child psychologist (IK), administered the Bayley-III in the current trial. Standardization exercises was conducted prior to the start of the study assessment until satisfactory level of agreement was reached. During the trial, 20% of the tests was double scored with an inter class correlation (ICC) ranging from 0.88 to 0.99 indicating high inter-rater agreement.

Background variables

Demographic information including education, income, pre-pregnancy and current weight, height, and nicotine use in pregnancy were collected through an electronic questionnaire at baseline in gestational week 18-19. Gestational length and birth weight were obtained from birth records by the mother's recall. Preterm delivery was defined as birth <gestational week 37 and low birthweight as <2500 g.

Iodine intake

Iodine intake was estimated from a structured 6-days iodine specific food diary designed and validated for this study (22). The food diary was filled out on six consecutive days at baseline and post-intervention (between gestational week 18-19 and gestational week 35-36 respectively) at the exact same days as the spot urinary samples. The food diary included food items of iodine rich foods (fish and seafood, milk and dairy products and eggs) in addition to supplements used. The food diary was developed specifically for this study and has been validated and described in details by Næss and Aakre et al. 2019 (22). The article includes information of iodine content of the specific food items used, including cod, for calculation of iodine intake. For data on the use of iodine supplements since becoming pregnant, we used data from an iodine specific food frequency questionnaire (FFQ), which is also validated and described in detailed by Næss and Aakre et al. 2019 (22). To retrieve the iodine content of the cod given in the intervention 30 individual cod samples from the batch used in the intervention were analyzed at the IMR using ICP-MS, the method has previously been described in detail elsewhere (23).

Statistical analysis

The power calculation for the sample size in the current study were based on data on a median (IQR) UIC of ~ 80 (80) $\mu\text{g/L}$ in pregnant women from the 'Little in Norway' cohort (9, 24). The sample size was calculated to detect a 30% difference in UIC between the intervention and control groups. The intended sample size was thus 60 individuals per group (total of 120) (0.05 1-tailed alpha, power 0.952). To account for attrition, twenty-four (20%) subjects were added reaching a final intended sample size of 144 (18, 25).

The continuous variables are summarized as mean \pm SD or median (interquartile range (IQR)) and the categorical variables are described in frequency and percent. Analyses were performed on an intention to treat basis, and missing cases were omitted from the dataset (listwise deletion) (26). Normality was assessed by testing the distribution of continuous variables against a normal distribution using the Shapiro Wilk W test. For the primary outcome, we present the median (IQR) UIC $\mu\text{g/L}$ at baseline and post intervention. UIC data were transformed using \log_{10} to correct for positive skewness. In the main primary analysis, the student t-test was used to compare potential within and between group differences, and a one-way ANCOVA was used to compare the differences in post intervention UIC $\mu\text{g/L}$ between the groups with baseline UIC $\mu\text{g/L}$ included as a covariate. We included baseline UIC in the model as it is believed that the post intervention UIC to some degree depend on the baseline UIC $\mu\text{g/L}$. We also assessed whether UIC had remained stable, increased or decreased between the baseline and post intervention. An increase and decrease were defined as a change larger than $\pm 10\%$ from baseline to post intervention, which were chosen according to the measurement uncertainty of 20%. The student t-test was used to compare potential within and between group differences in thyroid hormone levels (TSH, fT3 and fT4), and a one-way ANCOVA was used to compare the differences in post intervention concentrations between the groups with the baseline concentrations included as a covariate. For the secondary outcome, we present the mean (SD) of the Bayley-III cognitive composite score, the language scaled scores (receptive and expressive) and composite scores and the motor scaled scores (fine and gross) and composite scores. We used a student t-test to compare the means between the groups.

Statistical analyses were performed using Statistical Package for the Social Sciences (SPSS® Statistics Version 25).

Results

A total of 165 pregnant women showed interest in participating in the study by contacting the study secretariat. From January 2016 until February 2017 a total of 137 pregnant women were enrolled in the study and signed the informed consent. Between the enrolment and randomization, four participants withdrew from the study, and in total 133 pregnant women were randomized to the intervention group ($n=68$) or the control group ($n=65$). Between randomization and post-testing, nine participants dropped out of the study (six (8.8%) in the intervention group and three (4.6%) in the control group). In addition, two participants were lost to follow up due to preterm delivery. Hence, for the primary outcome (UIC), 122 participants [$n=61$ (89.7%) in the intervention group and $n=61$ (93.8%) in the control group] were included in analysis. For the secondary outcome (Bayley-III), 112 participants [$n=57$ (83.8%) in the intervention group and $n=55$ (84.6%) in the control group] were included in the analyses. There were no differences in any baseline characteristics between participants who withdrew from the study and those who completed the study at 11 months (see Supplementary Table 1). An overview of the recruitment and flow of participants through the trial is outlined in **Figure 1**. Baseline characteristics for the randomized pregnant women were similar in the two groups (**Table 1**). Mean compliance score was 77, ranging from 35 to 102. More than 70% of the participants had a compliance score of more than 70. Approximately 50% of the participants had a compliance score of more than 80. Less than 10% had a compliance score of <50. The mean (SD), median (IQR), and 5 percentile intakes of the received cod in the intervention group was 306 (62), 318 (275-356), and 175 g per week, respectively. The mean (IQR, min, max) analyzed iodine content of the cod given in the intervention based on analysis of 30 individual cod samples was 81 (46-71, 31, 630) $\mu\text{g}/100\text{g}$. There were no reported adverse events during or after the intervention.

In total, six infants were delivered preterm (< gestational week 37) ($n=3$ in the control group, $n=3$ in the intervention group), and three infants were born with low

birthweight (<2500 g) ($n=2$ in the control group, $n=1$ in the intervention group). There was no difference between gestational length (control group: mean (SD): 40.2 (2.6) weeks, intervention group: mean (SD): 40.3 (1.8) weeks) and birth weight (control group: mean (SD): 3442 (589) g, intervention group: mean (SD): 3541 (478) g) between the groups.

For the main analysis, the UIC was significantly higher in the intervention group post intervention (median (IQR) 98 (64-145) $\mu\text{g/L}$), compared to the control group (median (IQR) 73 (52-120) $\mu\text{g/L}$) ($p=0.028$). The difference between the groups was still significant after adjusting for baseline UIC $\mu\text{g/L}$ (log mean (SD) control group: 1.89 (0.027), log mean (SD) intervention group: 1.96 (0.027), mean difference between log means: 0.076 (95% CI 0.001, 0.150), $p=0.048$) (**Table 2**). The estimated iodine intake post intervention was significantly higher ($p=0.001$) in the intervention group (median (IQR, min, max) 218 (156-323, 45,481), $\mu\text{g/day}$, $n=61$) compared to the control group (median (IQR, min, max) 146 (87-264, 32-423), $\mu\text{g/day}$, $n=61$).

For the secondary outcome, the intervention group had a significant lower cognitive composite score on the Bayley-III compared to the control group, when the infants were 11 months of age (log mean (SD) control group: 1.99 (0.049), log mean (SD) intervention group: 1.97 (0.04), mean difference between log means: 0.016 (95% CI 0.0004, 0.0321), $p=0.045$). There were no significant differences in the Bayley-III language- or motor composite scores between the groups (**Table 3**).

Table 4 shows change in UIC from baseline to post intervention in the control and intervention groups. The number of participants with an increase in UIC was 28% in the control group and 36% in the intervention group, while the number of participants with a decrease in UIC was 51% and 36% in the control and intervention group, respectively. The thyroid hormones TSH, fT3 and fT4 did not differ between the groups post intervention (**Table 5**). There was no difference in DHA status neither at baseline nor post intervention between the groups (see Supplementary Table 2).

Discussion

To our knowledge, this is the first RCT to investigate the effect of nonfortified iodine rich food consumption in pregnancy on maternal iodine status and infant neurodevelopment. In concordance with other recent studies in Norwegian and other European pregnant populations, the pregnant women in this study were mildly-to-moderately iodine deficient at baseline. After the intervention with two meals of cod for a period of 16 weeks in pregnancy (GW 20-36), median UIC was significantly higher in the intervention group compared to the control group. However, the median UIC values remained below the recommended UIC of 150 $\mu\text{g/L}$. The children's language and motor scores on the Bayley-III were similar in both groups when assessed at 11 months of age while the cognitive score was better in the control group.

Our results suggest that it is possible to improve iodine status during pregnancy using a nutrition-sensitive food system approach to meet the dietary requirements of a population (27). Several RCT's have previously been conducted to improve iodine status in pregnancy, using iodine containing supplements (28). In countries without iodized salt, as in Norway and the UK, it is important to include one or several key sources of iodine in the habitual diet in order to achieve an adequate iodine intake. The present study shows that cod has a potential as an important dietary source of iodine in pregnancy when included in the diet. However, median UIC after the intervention did not meet the epidemiological criteria for adequate iodine nutrition (median UIC between 150-250 $\mu\text{g/L}$) suggested by the WHO (2). Though, the suggested limit of ~ 100 $\mu\text{g/L}$, based on the Nordic Nutrition Recommendations were almost met with a median of 98 $\mu\text{g/L}$ in the intervention group (5).

The intervention group had a significantly lower score on the cognitive composite score than the control group, while the language and motor scores were similar between the groups. Since this is the first RCT aiming at increasing iodine status in pregnancy through a food-based approach, and subsequently assessing child development, comparable studies are lacking. In regions of severe iodine deficiency, iodine supplementation in pregnancy have been shown to reduce the incidence of cretinism and

improve motor development in children (16). However, the effects of iodine supplementation during pregnancy in mildly-to-moderately iodine deficient populations are still unclear (16). A Cochrane review from 2017, including 11 trials, concluded that there was insufficient evidence for either the benefit or harm of iodine supplementation during pregnancy on child neurodevelopment due to small participant numbers and low quality of trials (28). Only two of the RCTs included developmental outcomes in children and in those, there were no difference in the developmental outcomes between the groups (29, 30). The first randomized, double-blinded, placebo-controlled trial in mildly-to-moderately iodine deficient pregnant women was conducted from 2008 to 2011 in India and Thailand. They found no effect of 200 µg iodine per day from early pregnancy (<gestational week 14) until delivery on child cognition at 5-6 years of age (31). It is important to note that albeit the median UIC of the two cohorts was indicative of mild deficiency at baseline, one of the cohorts were actually iodine sufficient, which potentially could have confounded the results. However, an analysis of secondary outcomes at age one year showed higher scores for expressive language measured by the Bayley III in the placebo group than in the iodine intervention group.(31). Likewise, we found a higher Bayley III score on cognition in the control group than in the intervention group. However, the effect size of the difference between the groups were small, and mean cognitive level was close to average for both the control and intervention group. Still, the mean difference in Bayley-III cognitive score of 4 points could reflect important differences in learning and development. Follow-up studies investigating stability and trajectories are needed to conclude on the impact of these differences. Although more uncertain on an individual level, small effects may be meaningful on a public health level. Results from a few RCT's, and observational studies suggest that introducing an iodine supplement after the onset of pregnancy in women with mild-to-moderate iodine deficiency can have adverse effect on thyroid hormones and infant development (28, 32-35). Furthermore, a recent systematic review and meta-analysis of the effects of iodine supplementation on thyroid function and child neurodevelopment in mildly-to-moderately iodine-deficient pregnant women concluded that there is insufficient good-quality evidence to support current recommendations for iodine supplementation in pregnancy in areas of mild-to-

moderate deficiency (36). Despite these ambiguous results, routine iodine supplementation in pregnancy is recommended by health authorities across the world.

The possible underlying mechanisms of all iodine deficiency disorders (IDDs), including impaired child development, are inadequate thyroid hormone production and their disturbed action in target tissues (37). Correcting iodine deficiency has proven beneficial in order to prevent IDDs worldwide (38, 39). However, iodine supplementation or fortification has also been associated with increased incidence of thyroid dysfunction (40). Even though we managed to increase the iodine status in the intervention group, it might have been too late as the first trimester of pregnancy is crucial as the fetal brain development is dependent on maternal thyroid hormone transfer (41). Moreover, not all participants in the intervention group had an increase in UIC from baseline to post intervention, which may have interfered with the secondary outcome. Still, for those participants who had a UIC <50 µg/L at baseline, the intervention group differed from the control group in that all the participants had an increase in UIC from baseline to post intervention. Nevertheless, we cannot exclude the possibility that the alterations in iodine intake during pregnancy could have a negative effect on maternal thyroid function and hence child cognition. Noteworthy, a temporal stunning of thyroid function has been discussed to be as a consequence of changes in iodine intake in mildly-to-moderately iodine deficient populations, and not solely as a consequence of excessive iodine intake (42, 43). In the current study, the women's iodine intake could not be characterized as excessive.

The differences in UIC concentration and estimated iodine intake between groups after the intervention were not reflected in the thyroid hormones that were similar in the groups post-intervention. However, individuals have a genetic set-point for thyroid hormone concentrations and, despite wide inter-individual variation, there is a low index of individuality (44). Nonetheless, there is a possibility that a small change in the habitual maternal iodine intake during pregnancy, rather than prior to pregnancy, might have changed fetal thyroid hormone status during this vulnerable period of neurodevelopment. Thus, our finding urges further attention.

A key limitation in this study could be the onset of the intervention as timing of exposure to maternal iodine insufficiency is a modulator for its effect on outcomes (45). In the attempt of a non-biased source population, the women were recruited before their first meeting with the public health care system at gestational week 18. Thus, baseline testing, randomization and onset of the dietary intervention did not commence before gestational week 18 and 19. Blinding is a challenge in RCTs with food. We have previously conducted RCTs with fish in kindergarten children and youths preparing identical meals to intervention and control group, the only difference being the protein source (46, 47). However, due to both texture and taste, blinding was not possible. RCTs with single dietary components are inconclusive in confirming health outcomes from observational studies. This could be due to the complexity of food containing several bioactive components. Thus, RCT's with food are still valuable despite the challenge of blinding (48). A limitation, or rather, a consequence of a dietary intervention introducing a food (e.g. lean fish) twice a week is replacement of another habitual food (e.g. fatty fish). While lean fish is a good source of iodine, fatty fish is an excellent source of DHA, a nutrient also playing a critical role in brain development (49, 50). Details regarding the seafood intake of the participants in this study has recently been described in detail elsewhere (51). While the control group had a stable intake of both lean and fatty fish, the intervention group increased their intake of lean fish and decreased their intake of fatty fish from baseline to post intervention. The intervention group had a higher DHA status compared to the control group post-intervention, although, the difference was not significant. Still, the higher DHA status in the intervention group is not likely to confound the effects of interest or have a large impact on the cognitive outcomes. As lean fish also is a source of mercury, we also measured mercury status at baseline and post intervention (51). However, the concentrations of mercury were low and only a small difference between the groups were observed, so we do not believe this has affected the affected the results. Furthermore, the iodine content of the cod given in the intervention was low compared to reported values in the Norwegian food composition table. Laboratory analysis also showed large variation in the individual cod fillets. Thus, the dose given in each meal and to each participant in the intervention group is believed to have varied, introducing a dose bias. We did not correct for multiple testing for the secondary outcome, and thus the risk of detecting false

positive results (type 1 error) might have increased. However, correction for multiple testing has been debated as it also decreases the statistical power and increases the risk of not detecting real differences (type 2 error) (52).

Strengths of this study is the high compliance, and the analysis of a pooled sample of six spot urine samples at baseline and post intervention in both groups including both weekdays and weekend. Although, ten samples are suggested for assessing individual iodine status, six samples could be enough to categorize individual status based on UIC as there was a strong agreement with the food diary, recorded on the same six consecutive days (22, 53).

In conclusion, an increased intake of iodine through consumption of cod in mildly-to-moderately iodine deficient pregnant women resulted in an increased iodine status measured as UIC. The intervention had no measurable effect on maternal thyroid function, and the infants of the mothers in the intervention group had a lower cognitive score compared to the control group. A follow up of these children over time will indicate if these differences persist. While the overall literature is ambiguous, there is a need to investigate further both positive and possible adverse effects of increasing iodine intake in pregnancy, rather than ideally before conception, to assure the safety of the present guidelines.

Declarations

Acknowledgments

The authors are very grateful for all the women who participated in the study. The authors are thankful for all technical and laboratory assistance, especially to Elisabeth R. Ødegaard and technicians at the Inorganic Laboratory at IMR. The Mommy's Food study is financially supported by The Norwegian Seafood Research Fund (<http://www.fhf.no/>). The funder had no role in design or the decision to publish this protocol. The cod used in the trial was purchased, after tender, from Lerøy A/S, Bergen, Norway.

Author Disclosure Statement

The authors declare they have no actual or potential competing financial interest of receiving the grant to perform this study, or no other actual or potential competing financial interests exist.

Ethics approval and consent to participate

The Regional Committee for Medical and Health Research Ethics West has granted ethical approval for the original randomized controlled Mommy's Food trial (ref:2015/879). Written informed consent from the mother and assent from the child were obtained. The trial is registered in ClinicalTrials.gov (NCT02610959).

Availability of data materials

Requests for data collected in the Mommy's Food study (such as deidentified participant data) can be made to the corresponding author following publication, and requests will be considered on an individual basis. Any requests require completion and approval of the application for use of data from the Mommy's Food study. The trial project group will review and, if acceptable and approved by the Regional Committee for Medical and Health Research Ethics West, Norway, provide approval of the request. A signed data sharing access agreement will be required. The data will be provided as a SPSS dataset. Any other format requests might incur costs to the requestor. To facilitate the data access process please contact. Data will be available after publication of the study results. Maria Wik Markhus at maria.wik.markhus@hi.no and mammasmat@hi.no.

Funding

The Norwegian Seafood Research Fund (<http://www.fhf.no/>) funded this study. The funder of the study had no role in study design, data collection, data analysis, data interpretation, or writing of the report. The corresponding author had full access to all the data in the study and had the final responsibility for the decision to submit for publication. The study was not overseen by a data monitoring committee.

Authors' contributions

MK, LD, MWM, MH, IK designed the study, LKM, IN, MWM, LD, MK, MH and IK conducted the study, MWM, SN and IA did the statistical analysis, MWM wrote the first drafts of the report with input from MH, IA, SN, IK and MK. All authors critically reviewed the final manuscript.

References

1. Andersson M, Karumbunathan V, Zimmermann MB 2012 Global iodine status in 2011 and trends over the past decade. *The Journal of nutrition* **142**:744-750.
2. WHO, UNICEF, ICCIDD. 2007 Assessment of iodine deficiency disorders and monitoring their elimination. World Health Organisation.
3. Zimmermann MB, Gizak M, Abbott K, Andersson M, Lazarus JH 2015 Iodine deficiency in pregnant women in Europe. *The lancet Diabetes & endocrinology* **3**:672-674.
4. Zimmermann MB 2008 Iodine deficiency in pregnancy and the effects of maternal iodine supplementation on the offspring: a review. *The American journal of clinical nutrition* **89**:668S-672S.
5. Nordic Council of Ministers 2014 Nordic Nutrition Recommendations 2012: integrating nutrition and physical activity. Nordic Council of Ministers.
6. Abel MH, Korevaar TI, Erlund I, Villanger GD, Caspersen IH, Arohonka P, Alexander J, Meltzer HM, Brantsæter AL 2018 Iodine intake is associated with thyroid function in mild to moderately iodine deficient pregnant women. *Thyroid* **28**:1359-1371.
7. Markhus M, Dahl L, Moe V, Abel M, Brantsæter A, Øyen J, Meltzer H, Stormark K, Graff I, Smith L 2018 Maternal iodine status is associated with offspring language skills in infancy and toddlerhood. *Nutrients* **10**:1270.
8. Brantsæter AL, Abel M, Haugen M, Meltzer HM 2013 Risk of suboptimal iodine intake in pregnant Norwegian women. *Nutrients* **5**:424-440.
9. Dahl L, Markhus MW, Sanchez P, Moe V, Smith L, Meltzer HM, Kjellevoid M 2018 Iodine Deficiency in a Study Population of Norwegian Pregnant Women -Results from the Little in Norway Study (LiN). *Nutrients* **10**:513.
10. Henjum S, Aakre I, Lilleengen A, Garnweidner-Holme L, Borthne S, Pajalic Z, Blix E, Gjengedal E, Brantsæter AL 2018 Suboptimal iodine status among pregnant women in the Oslo area, Norway. *Nutrients* **10**:280.
11. Berg V, Nøst TH, Skeie G, Thomassen Y, Berlinger B, Veyhe AS, Jorde R, Odland JØ, Hansen S 2017 Thyroid homeostasis in mother–child pairs in relation to maternal iodine status: The misa study. *European journal of clinical nutrition* **71**:1002.

12. The Norwegian Directorate of Health 2019 Utviklingen i norsk kosthold 2018 [Trends in the Norwegian diet]. The Norwegian Directorate of Health, Oslo.
13. Carlsen M, Andersen L, Dahl L, Norberg N, Hjartåker A 2018 New Iodine Food Composition Database and Updated Calculations of Iodine Intake among Norwegians. *Nutrients* **10**:930.
14. Pharoah P, Butfield I, Hetzel B 1971 Neurological damage to the fetus resulting from severe iodine deficiency during pregnancy. *The Lancet* **297**:308-310.
15. WHO U 2007 Reaching optimal iodine nutrition in pregnant and lactating women and young children. Geneva, Switzerland: World Health Organization.
16. Zhou SJ, Anderson AJ, Gibson RA, Makrides M 2013 Effect of iodine supplementation in pregnancy on child development and other clinical outcomes: a systematic review of randomized controlled trials. *American Journal of Clinical Nutrition* **98**:1241-1254.
17. Pearce EN, Lazarus JH, Moreno-Reyes R, Zimmermann MB 2016 Consequences of iodine deficiency and excess in pregnant women: an overview of current knowns and unknowns. *Am J Clin Nutr* **104 Suppl 3**:918S-923S.
18. Markhus MW, Kvestad I, Midtbø LK, Nerhus I, Ødegaard ER, Graff IE, Lie Ø, Dahl L, Hysing M, Kjellevoid M 2018 Effects of cod intake in pregnancy on iodine nutrition and infant development: study protocol for Mommy's Food-a randomized controlled trial. *BMC nutrition* **4**:7.
19. Araujo P, Nguyen T-T, Frøyland L, Wang J, Kang JX 2008 Evaluation of a rapid method for the quantitative analysis of fatty acids in various matrices. *Journal of chromatography A* **1212**:106-113.
20. Bayley N 2006 Bayley scales of infant and toddler development. PsychCorp, Pearson.
21. Bayley N 2009 Bayley Scales of Infant and Toddler Development (Bayley-III) Norsk manualsupplement. Third Edition ed, San Antonio.
22. Næss S, Aakre I, Kjellevoid M, Dahl L, Nerhus I, Midtbø LK, Markhus MW 2019 Validation and reproducibility of a new iodine specific food frequency questionnaire for assessing iodine intake in Norwegian pregnant women. *Nutrition journal* **18**:62.

23. Nerhus I, Markhus MW, Nilsen BM, Øyen J, Maage A, Ødegård ER, Midtbø LK, Frantzen S, Kögel T, Graff IE 2018 Iodine content of six fish species, Norwegian dairy products and hen's egg. *Food & nutrition research* **62**.
24. Moe V, Fredriksen E, Kjellefold M, Dahl L, Markhus MW, Stormark KM, von Soest T, Olafsen KS, Vannebo UT, Smith L 2019 Little in Norway: a prospective longitudinal community-based cohort from pregnancy to child age 18 months. *BMJ Open* **9**:e031050.
25. Faul F, Erdfelder E, Lang A-G, Buchner A 2007 G* Power 3: A flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behavior research methods* **39**:175-191.
26. Nich C, Carroll KM 2002 'Intention-to-treat' meets 'missing data': implications of alternate strategies for analyzing clinical trials data. *Drug and alcohol dependence* **68**:121-130.
27. Uccello E, Kauffmann D, Calo M, Streissel M 2017 Nutrition-sensitive agriculture and food systems in practice. *FAO*.
28. Harding KB, Peña-Rosas JP, Webster AC, Yap CM, Payne BA, Ota E, De-Regil LM 2017 Iodine supplementation for women during the preconception, pregnancy and postpartum period. *Cochrane Database of Systematic Reviews* **3**:CD011761.
29. Brucker-Davis F, Ganier-Chauliac F, Gal J, Panaïa-Ferrari P, Pacini P, Fénelichel P, Hiéronimus S 2015 Neurotoxicant exposure during pregnancy is a confounder for assessment of iodine supplementation on neurodevelopment outcome. *Neurotoxicology and teratology* **51**:45-51.
30. Zhou SJ, Skeaff SA, Ryan P, Doyle LW, Anderson PJ, Kornman L, Mcphee AJ, Yelland LN, Makrides M 2015 The effect of iodine supplementation in pregnancy on early childhood neurodevelopment and clinical outcomes: results of an aborted randomised placebo-controlled trial. *Trials* **16**:563.
31. Gowachirapant S, Jaiswal N, Melse-Boonstra A, Galetti V, Stinca S, Mackenzie I, Thomas S, Thomas T, Winichagoon P, Srinivasan K, Zimmermann MB 2017 Effect of iodine supplementation in pregnant women on child neurodevelopment: a randomised, double-blind, placebo-controlled trial. *Lancet Diabetes Endocrinol* **5**:853-863.

32. Murcia M, Rebagliato M, Iniguez C, Lopez-Espinosa M-J, Estarlich M, Plaza B, Barona-Vilar C, Espada M, Vioque J, Ballester F 2011 Effect of iodine supplementation during pregnancy on infant neurodevelopment at 1 year of age. *American journal of epidemiology* **173**:804-812.
33. Rebagliato M, Murcia M, Álvarez-Pedrerol M, Espada M, Fernández-Somoano A, Lertxundi N, Navarrete-Muñoz E-M, Fornis J, Aranbarri A, Llop S, Julvez J, Tardón A, Ballester F 2013 Iodine Supplementation During Pregnancy and Infant Neuropsychological Development: INMA Mother and Child Cohort Study. *American Journal of Epidemiology* **177**:944-953.
34. Abel MH, Caspersen IH, Meltzer HM, Haugen M, Brandlistuen RE, Aase H, Alexander J, Torheim LE, Brantsaeter AL 2017 Suboptimal Maternal Iodine Intake Is Associated with Impaired Child Neurodevelopment at 3 Years of Age in the Norwegian Mother and Child Cohort Study. *Journal of Nutrition* **147**:1314-1324.
35. Abel MH, Brandlistuen RE, Caspersen IH, Aase H, Torheim LE, Meltzer HM, Brantsaeter AL 2018 Language delay and poorer school performance in children of mothers with inadequate iodine intake in pregnancy: results from follow-up at 8 years in the Norwegian Mother and Child Cohort Study. *European Journal of Nutrition* **58**:3047–3058.
36. Dineva M, Fishpool H, Rayman MP, Mendis J, Bath SC 2020 Systematic review and meta-analysis of the effects of iodine supplementation on thyroid function and child neurodevelopment in mildly-to-moderately iodine-deficient pregnant women. *The American Journal of Clinical Nutrition*.
37. Eastman CJ, Zimmermann MB 2018 The iodine deficiency disorders Endotext [Internet]. Vol. MDText. com, Inc.
38. Pearce EN, Andersson M, Zimmermann MB 2013 Global iodine nutrition: where do we stand in 2013? *Thyroid* **23**:523-528.
39. Markou K, Georgopoulos N, Kyriazopoulou V, Vagenakis A 2001 Iodine-induced hypothyroidism. *Thyroid* **11**:501-510.

40. Laurberg P, Jørgensen T, Perrild H, Ovesen L, Knudsen N, Pedersen IBI, Rasmussen LB, Carlé A, Vejbjerg P 2006 The Danish investigation on iodine intake and thyroid disease, DanThyr: status and perspectives. *European Journal of Endocrinology* **155**:219-228.
41. Bernal J, Nunez JJ 1995 Thyroid hormones and brain development. *European Journal of Endocrinology* **133**:390-398.
42. Zimmermann MB 2008 Iodine requirements and the risks and benefits of correcting iodine deficiency in populations. *Journal of trace elements in medicine and biology* **22**:81-92.
43. Delange F 1994 The disorders induced by iodine deficiency. *Thyroid* **4**:107-128.
44. Waise A, Price HC 2009 The upper limit of the reference range for thyroid-stimulating hormone should not be confused with a cut-off to define subclinical hypothyroidism. *Annals of clinical biochemistry* **46**:93-98.
45. Zoeller R, Rovet J 2004 Timing of thyroid hormone action in the developing brain: clinical observations and experimental findings. *Journal of neuroendocrinology* **16**:809-818.
46. Øyen J, Kvestad I, Midtbø LK, Graff IE, Hysing M, Stormark KM, Markhus MW, Baste V, Frøyland L, Koletzko B 2018 Fatty fish intake and cognitive function: FINS-KIDS, a randomized controlled trial in preschool children. *BMC medicine* **16**:41.
47. Handeland K, Skotheim S, Baste V, Graff IE, Frøyland L, Lie Ø, Kjellevoid M, Markhus MW, Stormark KM, Øyen J 2018 The effects of fatty fish intake on adolescents' nutritional status and associations with attention performance: results from the FINS-TEENS randomized controlled trial. *Nutrition journal* **17**:30.
48. Jacobs DR, Tapsell LC 2007 Food, not nutrients, is the fundamental unit in nutrition. *Nutrition reviews* **65**:439-450.
49. Lauritzen L, Brambilla P, Mazzocchi A, Harsløf L, Ciappolino V, Agostoni C 2016 DHA effects in brain development and function. *Nutrients* **8**:6.
50. Innis SM 2007 Dietary (n-3) fatty acids and brain development. *The Journal of nutrition* **137**:855-859.

51. Næss S, Kjellevold M, Dahl L, Nerhus I, Midtbø LK, Bank MS, Rasinger JD, Markhus MW 2020 Effects of seafood consumption on mercury exposure in Norwegian pregnant women: A randomized controlled trial. *Environment International* **141**:105759.
52. Rothman KJ 2014 Six persistent research misconceptions. *Journal of general internal medicine* **29**:1060-1064.
53. König F, Andersson M, Hotz K, Aeberli I, Zimmermann MB 2011 Ten repeat collections for urinary iodine from spot samples or 24-hour samples are needed to reliably estimate individual iodine status in women. *The Journal of nutrition* **141**:2049-2054.

Table 1. Baseline characteristics of study participants within each group in Mommy's Food trial.

Characteristic	Control		Intervention	
	<i>n</i>		<i>n</i>	
Age, in years. mean (SD)	65	29.1 (3.5)	68	29.6 (4.0)
BMI, kg/m ²	62	23.3 (4.3)	68	22.9 (3.9)
Education, %	63		68	
≥ 12 years	7	11.1	11	16.2
13-16 years	15	23.8	18	26.5
> 16 years	41	65.1	39	57.4
Household Income (NOK ^a), %	63		68	
Low (< 200,000-549,000)	15	23.8	23	33.8
Medium (550,000-1,249,999)	40	63.5	36	52.9
High (1,250,000- >2,000,000)	8	12.7	9	13.2
Nicotine use in pregnancy ^b , yes (n) %	62		68	
≤ gestational week 8	7	11.3	5	7.4
> gestational week 8	0	0	0	0
Iodine intake (µg/day), median (IQR) ^c	65	133 (81-240)	68	152 (92-267)
Milk and Dairy	65	67 (43-92)	68	62 (22-76)
Egg	65	7 (3-14)	68	7 (3-14)
Seafood ^b	65	33 (94)	68	42 (94)
Supplements (users only)	21	175 (135-200)	25	150 (131-175)
Iodine supplement use since becoming pregnant, n (%) ^d	59	21 (32)	63	29 (43)

^a 100 NOK= approximately 11.6 USD/10.2 EUR, ^b No participants reported use of nicotine after gestational week 8. Abbreviations: BMI, body mass index, ^c Intake estimated from 6-day food diary at baseline, ^d Reported from food frequency questionnaire.

Table 2. Urinary iodine concentration at baseline and post intervention in Mommy's Food trial participants.

	Urinary iodine concentration ($\mu\text{g/L}$) ^a		Difference between groups post intervention	
	Baseline	Post	Crude	Adjusted
	Median (IQR)	Median (IQR)	<i>p-value</i> ^b	<i>p-value</i> ^c
Control (n= 61)	85 (55-130)	73 (52-120)		
Intervention (n= 61)	88 (64-130)	98 (64-145)	<i>0.028</i>	<i>0.048^d</i>

^a Analysed in a pooled sample of six spot samples collected on six consecutive days at each timepoint. ^b Independent sampled *t*-test for comparison of log transformed values. ^c One-way analysis of covariance (ANCOVA) for comparison of differences between control and intervention group adjusted for baseline urinary iodine concentration ($\mu\text{g/L}$). ^d $\eta^2=0.033$.

Table 3. Infant neurodevelopment assessed at 11 months of age in Mommy's Food trial participants.

	Control (n=55)	Intervention (n=57)	
Neurodevelopment scores ^a	Mean (SD)	Mean (SD)	p-value ^b
Cognitive composite score	99 (10)	95 (9)	0.045 ^c
Language composite scores	95 (8)	96 (8)	0.67
<i>Receptive language scaled score</i>	8 (2)	8 (2)	0.20
<i>Expressive language scaled score</i>	11 (1)	11 (1)	0.36
Motor composite scores	94 (8)	92 (7)	0.24
<i>Gross motor scaled score</i>	9 (2)	9 (2)	0.85
<i>Fine motor scaled score</i>	9 (1)	9 (2)	0.13

^a Neurodevelopment was assessed by the cognitive, language and motor scales of the Bayley Scales of Infant and Toddler Developmental 3rd edition. ^b Independent *t*-test of differences for comparison of log transformed values between groups. Language composite scores owing to normality. ^c Cohen's *d*= 0.42

Table 4. Change in urinary iodine concentration from baseline to post intervention in Mommy's Food trial participants.

UIC Baseline	UIC Post intervention						<i>p-value</i> ^c
	Control			Intervention			
	Increase n (%) ^a	Decrease n (%) ^a	Stable n (%) ^a	Increase n (%) ^b	Decrease n (%) ^b	Stable n (%) ^b	
<50 µg/L	4 (7)	5 (8)	3 (5)	7 (12)	0	0	0.034
50-100 µg/L	9 (15)	9 (15)	4 (7)	15 (25)	7 (12)	9 (15)	0.243
>100 µg/L	4 (7)	17 (28)	6 (10)	3 (5)	15 (25)	5 (8)	0.847
All	17 (28)	31 (51)	13 (21)	25 (41)	22 (36)	14 (23)	0.081

Increase and decrease was defined as a change larger than $\pm 10\%$ from baseline to post intervention. ^a Percent within control group. ^b Percent within intervention group. ^c

Differences tested between increase and decrease in the control and intervention group using Pearson's chi square test or Fisher's exact test.

Table 5. TSH, fT3 and fT4 in intervention and control group at baseline (gw 18) and post intervention (gw 36) in Mommy's Food trial participants.

	Thyroid hormones		Difference between groups post intervention	
	Baseline	Post	Crude	Adjusted
	Mean (SD)	Mean (SD)	<i>p-value</i> ^a	<i>p-value</i> ^b
	TSH (mIU/L)			
Control (<i>n</i> = 58)	1.6 (0.7)	1.8 (0.8)	<i>0.94</i>	<i>0.89</i>
Intervention (<i>n</i> = 61)	1.6 (0.8)	1.8 (0.8)		
	fT3 (pmol/L)			
Control (<i>n</i> = 58)	4.3 (0.4)	3.8 (0.4)	<i>0.18</i>	<i>0.10</i>
Intervention (<i>n</i> = 61)	4.2 (0.5)	3.9 (0.4)		
	fT4 (pmol/L)			
Control (<i>n</i> = 58)	13.8 (1.5)	13.5 (1.9)	<i>0.87</i>	<i>0.60</i>
Intervention (<i>n</i> = 61)	14.0 (1.7)	13.4 (1.4)		

^a Independent *t*-test of differences between groups post intervention. ^b One-way analysis of covariance (ANCOVA) for comparison of differences between control and intervention group adjusted for baseline levels. Abbreviations: gw, gestational week; TSH, thyroid stimulating hormone; fT3, free triiodothyronine; fT4, free thyroxine.

Figure legend




Figure 1. Trial profile depicting the flow of participants through the intervention trial. Abbreviations: UIC, urinary iodine concentration; Bayley III, Bayley Scales of Infant and Toddler Developmental 3rd edition.